



Liu, X.-J., Hu, H., & Chang, L. (2001). Fragmented condensate ground state of trapped weakly interacting bosons in two dimensions.

Originally published in *Physical Review Letters*, 87(3).
Available from: <http://dx.doi.org/10.1103/PhysRevLett.87.030404>.

Copyright © 2001 The American Physical Society.

This is the author's version of the work. It is posted here with the permission of the publisher for your personal use. No further distribution is permitted. If your library has a subscription to this journal, you may also be able to access the published version via the library catalogue.

The definitive version is available at <http://prl.aps.org/>.



Fragmented Condensate Ground State of Trapped Weakly Interacting Bosons in Two Dimensions

Xia-Ji Liu¹, Hui Hu¹, Lee Chang², Weiping Zhang¹, Shi-Qun Li^{1,3}, and Yu-Zhu Wang⁴

¹*Department of Physics, Tsinghua University, Beijing 100084, China*

²*Center for Advanced Study, Tsinghua University, Beijing 100084, China*

³*Center for Atomic and Molecular Nanosciences, Tsinghua University, Beijing 100084, China*

⁴*Shanghai Institute of Optics and Fine Mechanics, Chinese Academy of Sciences, Shanghai 201800, China*

The ground state and its structure for a rotating, harmonically trapped N -Boson system with a weak repulsive contact interaction are studied as the angular momentum L increases up to $3N$. We show that the ground state is generally a fragmented condensate due to angular momentum conservation. In response to an (arbitrarily weak) asymmetric perturbation of the trap, however, the fragmented ground state can be transformed into a single condensate state. We manifest this intrinsic instability by calculating the conditional probability distributions, which show patterns analogous to the boson density distributions predicted by mean-field theory.

PACS numbers:03.75.Fi, 05.30.Jp, 67.40.Db, 67.40.Vs

Following the experimental realization of a dilute atomic Bose-Einstein condensate (BEC), the formation and properties of vortices in an atomic BEC have caused considerable interest both experimentally [1] and theoretically [2–9] in the past few years. Although the recent demonstrations of vortex states by several different groups are in the Thomas-Fermi limit of strong interatomic interaction, a great deal of attention have also been attached to the nonvanishing angular momentum states of weakly interacting N -Boson systems in harmonic traps. In Ref. [3], Wilkin *et al.* considered the case of an attractive interaction and showed that the ground state is *uncondensed* and is an example of the "fragmented" condensate discussed by Nozière and Saint James [10]. Mottelson [4], Bertsch and Papenbrock [5] considered the lowest energy quantum states of a repulsively interacting Bose gas when $L \leq N$. Wilkin *et al.* have employed a composite boson/fermion picture to describe configurations beyond the one-vortex state [6]. A more tractable mean-field calculation performed by Butts and Rokhsar revealed a succession of transitions between stable vortex patterns of differing symmetries in the high angular momentum regime [7]. The connection between the mean-field theory (MFT) and exact diagonalization scheme has been studied by Jackson *et al.* for a special case of $L = 2N$ [8]. Finally, some analytical results have also been reported for the lowest energy states [9].

In this Letter we address the question of whether the ground state of a weakly interacting N -Boson system with a given angular momentum is what one would normally expect, i.e., a state with a single coherent Bose condensate, in which a mean-field approximation is valid. We propose that this is not the case and the ground state is generally a *fragmented* condensate in the presence of the weakly repulsive interatomic interaction except $L/N = 0$ or 1 in the thermodynamic limit. By evaluating the macroscopic eigenvalues of the single-particle (SP) density matrix, we determine the degree of condensation. The origin of fragmentation turns out to be a requirement of the conservation of angular momentum.

As a result, by turning on an (arbitrarily weak) asymmetric perturbation of the trap, the fragmented ground state can be easily deformed to a single condensate state [11]. This intrinsic instability can be manifested by the conditional probability distributions (CPDs) calculated for the ground state, which show patterns analogous to boson density distributions predicted by MFT. Note that the weakly interacting N -Boson system considered here are quite similar to the spin-1 Bose gas studied by Ho and Yip [12], in which the fragmentation originates from the spin conservation.

We start from the model Hamiltonian describing N bosons in a two-dimensional harmonic trap interacting via a weak contact interaction. The SP spectrum is usually expressed in terms of the angular momentum quantum number m and the radial quantum number n_r , by $E_{n_r, m} = (2n_r + |m| + 1)\hbar\omega$. In the ground state of the system all the bosons are in states with $n_r = 0$, and with m being zero or having the same sign as the total angular momentum. In the second quantized form, the Hamiltonian reads

$$\begin{aligned} \mathcal{H} &= \mathcal{H}_0 + \mathcal{V}, \\ \mathcal{H}_0 &= \hbar\omega \sum_j (j+1) \hat{a}_j^\dagger \hat{a}_j, \\ \mathcal{V} &= \frac{1}{2}g \sum_{i,j,k,l} V_{ijkl} \hat{a}_i^\dagger \hat{a}_j^\dagger \hat{a}_k \hat{a}_l, \end{aligned} \quad (1)$$

where \mathcal{H}_0 is the SP oscillator Hamiltonian and \mathcal{V} is the two-body interaction between bosons. In the perturbative regime of weak interactions, $Ng \ll \hbar\omega$. The operator \hat{a}_j and \hat{a}_j^\dagger annihilate and create one boson in the SP oscillator state $|j\rangle$ with energy $(j+1)\hbar\omega$ and angular momentum $j\hbar$, respectively, and obey the bosonic commutation rules. The contact interaction elements are given by $V_{ijkl} = \delta_{i+j, k+l} 2^{-(i+j)} (i+j)! / (i!j!k!l!)^{1/2}$ [9], and most of them are actually vanishing. For a given total angular momentum L and number of bosons N , we consider the Fock space spanned by states $|\alpha\rangle = |n_0, n_1, \dots, n_k\rangle$

with $\sum_j n_j = N$ and $\sum_j j n_j = L$. Here n_j denotes the occupation of the j th SP oscillator state $|j\rangle$. There is a huge degeneracy corresponding to many different ways of distributing L quanta of angular momentum among N atoms. Here we restrict ourselves in a *truncated* Fock space of $0 \leq j \leq j_{\max} = 6$ [8,13]. To obtain the energy spectra and the corresponding eigenstates, we set up the matrix elements in the Fock space basis, and subsequently diagonalize the matrix by using the Davidson algorithm [14].

A fragmented ground state.—First consider the SP density matrix in the form of

$$\rho(\mathbf{r}, \mathbf{r}') = \sum_{ij} \psi_i^*(\mathbf{r}) \rho_{ij} \psi_j(\mathbf{r}') \quad (2)$$

with $\psi_m(\mathbf{r}) = \langle \mathbf{r} | m \rangle$. In Ref. [15], Yang showed that the appearance of condensation is associated with the single macroscopic eigenvalue (i.e. of order N) of the density matrix $\rho(\mathbf{r}, \mathbf{r}')$ with the "condensate wave function" being the associated eigenvector, while the case of more than one macroscopic eigenvalue has been referred to as a "fragmented" condensate [10]. The most important difference between the single and fragmented condensate is the lack of phase coherence of the latter. To find the eigenvalues of the SP density matrix, we write [15]

$$\rho_{ij} = \text{Sp} \hat{a}_i \rho \hat{a}_j^\dagger, \quad (3)$$

where the trace runs over all the $N - 1$ boson states, and the density matrix $\rho = |\Psi_{GS}\rangle \langle \Psi_{GS}|$. It is readily seen that the eigenvalues are nothing but the occupation numbers of the SP oscillator state due to the conservation of the total angular momentum, namely $\rho_{ij} = \delta_{ij} n_j$. It is difficult to give an explicit expression for the occupation numbers n_j . In the case of $L = N$, Wilkin *et al.* find that in the limit of $N \rightarrow \infty$, to the order $O(1/N)$ [3],

$$n_0 = 1, \quad n_1 = N - 2, \quad \text{and} \quad n_2 = 1. \quad (4)$$

They therefore conclude that the N -Boson system is fully condensed into the one-vortex state in the thermodynamic limit. More detailed information can be obtained from the exact diagonalization calculations [5]. In figure 1, we show the L dependence of the occupation numbers n_j and their fluctuations $\Delta n_j = (\langle \hat{n}_j^2 \rangle - \langle \hat{n}_j \rangle^2)^{1/2}$ for $j = 0, 1, 2, 3, 4$ for a system of $N = 40$ bosons. When $L \leq N$ the occupation numbers evolve rather smoothly as the angular momentum increases [5], while for $L > N$ there are many kinks in the curves, reflecting the complexity of the ground states. The most prominent feature in the figure is that for a high angular momentum there are generally at least two *significant* occupation numbers. For instance, at $L = 70$, the system has two large occupation numbers: $n_j \approx 9$ and 23 for $j = 0$ and 2, respectively. Evidently the case gives a "fragmented" condensate. Although the present calculation is performed in the case of $N = 40$, the conclusion that a fragmented condensate ground state exists universally applies to the

trapped, weakly interacting and rotating N -Boson systems with an arbitrary N including the thermodynamic limit $N \rightarrow \infty$ [7].

To examine the validity of the above statement, we investigate the N -dependence of the number of significantly occupied SP states by computing the inverse participation ratio [16,5]:

$$I_C = \sum_j (n_j/N)^2. \quad (5)$$

The I_C is the first nontrivial moment of the distribution of occupation numbers among the different SP states (note that $\sum_j n_j/N = 1$ by normalization). Its inverse $1/I_C$ qualitatively measures the number of significantly occupied SP states. For example, I_C would be unity for a system that only had a single macroscopic occupied states; the maximum value of N is reached in the opposite extreme, when all the SP oscillator states of the system are equally occupied. Figure 2 shows a plot of $1/I_C$ as a function of angular momentum L for a system of $N = 30, 40, 50$ and 60 bosons. It is easy to see that in the regime of $L/N < 1.6$ the value of $1/I_C$ varies smoothly as L/N increases and shows little dependence on N . In particular, the variation of the peak height at $L/N \approx 1.6$ is less than 3% as N increases from 30 to 60 (not shown in the figure). For $L/N > 1.6$, some irregular small oscillations appear in the curves. However, the overall profile of $1/I_C$ is still nearly independent of N . These small oscillations are purely due to the finite N effect [17] and decay gradually with increasing N . One may expect them to vanish in the limit of $N \rightarrow \infty$. Therefore, we conclude that $1/I_C$ can be further used to *qualitatively* measure the number of *macroscopically* occupied SP states in the thermodynamic limit, or in other words, to determine whether the ground state is fragmented or not.

As shown in figure 2, there are two global minima (≈ 1) at $L/N = 0$ and $L/N = 1$, which can be well interpreted as a signature of single condensates. For other values of L/N (especially in the high angular momentum), however, $1/I_C$ is generally larger than 2. This clearly indicates the fragmented nature of the corresponding ground states. Another notable feature in figure 2 is that the overall profile of $1/I_C$ exhibits a valley around $L/N = 1.0, 1.8$, and 2.4. This is consistent with the broad peaks of n_j at $L \approx 40, 70$ and 90 for $j = 1, 2$ and 3, respectively, as shown in figure 1. The number fluctuations Δn_j are in the order of $O(n_j^{1/2})$ for these peaks, exhibiting a *local* characteristic of a single coherent condensate.

The intrinsic spontaneous symmetry breaking of a fragmented state.—Let us now consider the stability of such a fragmented ground state. In Ref. [11,12], the authors argued that the fragmented state is inherently unstable to the formation of a single condensate of well-defined phase. The essential idea is that even a weak perturbation that breaks the conservation laws will rapidly generate phase coherence, modifying the density matrix deterministically to give a unique macroscopically occupied

SP state. To support this point, we first show that the fragmented state and its corresponding single condensate state have the same energies in the limit of $N \rightarrow \infty$, up to the order of $O(gN)$. A similar conclusion has been reported by Jackson *et al.* for the special case of $L = 2N$ [8]. In figure 3, the interaction energy V_{int} in units of gN^2 is plotted as a function of L/N for a system of $N = 20, 40$, and 60 bosons. As N increases, V_{int} becomes closer in value to that of a single condensate V_{int}^{mf} , as predicted by the MFT [7]. The inset shows the energy difference $\Delta V_{int} = V_{int}^{mf} - V_{int}$ in units of gN . It is readily seen that all the ΔV_{int} with different N are approximately located on a *universal* curve. This strongly suggests that ΔV_{int} can be described by an approximate form

$$\Delta V_{int} = \alpha gN \ll \hbar\omega, \quad (6)$$

where in the thermodynamic limit the factor $\alpha \sim 1$ depends on L/N only and the inequality comes from our assumption of weak interaction. As a result, even a perturbation of order $O(1/N)$ can be enough to drive the fragmented state into a single condensate state. This fact clearly indicates that the fragmented state will spontaneously break whatever the fragmentation was permitted by cylindrical symmetry in the first place.

This result can be understood in another way by considering the conditional probability distributions (CPDs) [18] that give the density correlation among bosons. We define the CPD for finding one boson at \mathbf{r} given another at \mathbf{v}_0 as

$$\mathcal{P}(\mathbf{r} | \mathbf{v}_0) = \frac{\langle \Psi_{GS} | \sum_{i \neq j} \delta(\mathbf{r} - \mathbf{r}_i) \delta(\mathbf{v}_0 - \mathbf{r}_j) | \Psi_{GS} \rangle}{(N-1) \langle \Psi_{GS} | \sum_j \delta(\mathbf{v}_0 - \mathbf{r}_j) | \Psi_{GS} \rangle}. \quad (7)$$

Unlike the usual density distribution that is cylindrically symmetric under rotational invariant confinement, the CPD is asymmetric and reflects an *intrinsic* density distribution [19].

What will an inherently unstable fragmented state evolve into if a weak perturbation is switched on? One may expect that the system will rapidly change into a state having the same intrinsic density distribution as the fragmented state, and simultaneously generate phase coherence [12]. In view of this, the CPD gives the tendency of a system's evolution and can be regarded as a measurement of the possible spontaneous symmetry breaking.

In figure 4, we show the L dependence of the CPDs for a system of $N = 40$ bosons. As expected, we observe the successive vortex-like patterns of differing symmetries, which are in good qualitative agreement with the mean-field calculations [7]. Both of them show a gradual transition for the formation of one- (fig.4a) and two-vortex-like (fig.4b) states in contrast to the rapid appearance of the three-vortex-like state (fig. 4c) [7]. As mentioned above, we identify this similarity as a signal of spontaneous symmetry breaking of fragmented states.

On the other hand, one should not confuse CPDs with the "true" vortex patterns predicted by the MFT [7]. The

latter has phase coherence, which is not just well-defined in CPDs. Besides this, they have a different physical mechanism for the vortex emergence with the increasing L . For example, our results seem to show that the one- and two-vortex are produced at the center of the cloud of condensate, in apparent contradiction to the prediction of the MFT that the vortex enters the cloud from the low-density periphery. These differences may be resolved through the Josephson tunneling experiment suggested by Leggett and Sols [20]. Certainly, more accurate theoretical studies on the fragmented state are required.

In conclusion, we have studied the ground state of a weakly interacting N -Boson system with a given angular momentum. We propose that the ground state is generally a fragmented condensate state, which is rather fragile in response to a weak asymmetric perturbation. By calculating the corresponding CPDs, we manifest this intrinsic instability. A comparison with the mean-field results is also given.

We thank Y. Zhou, Y.-X. Miao, and C.-P. Sun for their stimulating discussions. X.-J. Liu was supported by the NSF-China.

-
- [1] M. R. Matthews *et al.*, Phys. Rev. Lett. **83**, 2498 (1999); K. W. Madison *et al.*, Phys. Rev. Lett. **84**, 806 (2000); R. Onofrio *et al.*, Phys. Rev. Lett. **85**, 2228 (2000).
 - [2] D. S. Rokhsar, Phys. Rev. Lett. **79**, 2164 (1997); D. L. Feder, C. W. Clark and B. I. Schneider, Phys. Rev. Lett. **82**, 4956 (1999); A. A. Svidzinsky and A. L. Fetter, Phys. Rev. Lett. **84**, 5919 (2000); K.-P. Marzlin, W. Zhang, and B. C. Sanders, Phys. Rev. A **62**, 013602 (2000).
 - [3] N. K. Wilkin, J. M. F. Gunn, and R. A. Smith, Phys. Rev. Lett. **80**, 2265 (1998).
 - [4] B. Mottelson, Phys. Rev. Lett. **83**, 2695 (1999).
 - [5] G. F. Bertsch and T. Papenbrock, Phys. Rev. Lett. **83**, 5412 (1999); Phys. Rev. A **63**, 023616 (2001).
 - [6] N. K. Wilkin and J. M. F. Gunn, Phys. Rev. Lett. **84**, 6 (2000).
 - [7] R. A. Butts and D. S. Rokhsar, Nature **397**, 327 (1999). G. M. Kavoulakis, B. Mottelson, and C. J. Pethick, Phys. Rev. A, **62**, 063605 (2000).
 - [8] A. D. Jackson, G. M. Kavoulakis, B. Mottelson, and S. M. Reimann, Phys. Rev. Lett. **86**, 945 (2001).
 - [9] A. D. Jackson and G. M. Kavoulakis, Phys. Rev. Lett. **85**, 2854 (2000); R. A. Smith and N. K. Wilkin, Phys. Rev. A **62**, 061602 (2000); Wen-Jui Huang, Phys. Rev. A **63**, 015602 (2001).
 - [10] P. Nozières and D. Saint James, J. Phys. (Paris) **43**, 1133 (1982).
 - [11] D. S. Rokhsar, eprint cond-mat/9812260 (1998).
 - [12] T.-L. Ho and S. K. Yip, Phys. Rev. Lett. **83**, 5412 (1999).
 - [13] In the mean-field calculation, the occupancy of SP states with $j > 6$ is very low for $L \leq 2N$ [7]. For $2N < L \leq 3N$, our numerical results with $N = 20$ show that $n_j/N <$

0.03 ($j > 6$), therefore including such states does not alter our results qualitatively.

- [14] E. R. Davidson, *J. Comput. Phys.* **17**, 87 (1975).
- [15] C. N. Yang, *Rev. Mod. Phys.* **34**, 694 (1962).
- [16] L. Kaplan, *Nonlinearity* **12**, R1 (1999).
- [17] As we shall see, the ground state has an *intrinsic* vortex-like density distribution, which has a 2-fold and 3-fold symmetry for $2.0 > L/N > 1.6$ and $2.8 > L/N > 2.0$, respectively. A naive interpretation of *certain* stable ground states with p -fold symmetry is that some bosons possess p quanta of angular momentum and others have zero angular momentum [6]. In these ground states, when L is increased by one unit, the system will be puzzled as to how to distribute the extra angular momentum, until additional $p - 1$ units are supplied. This causes an oscillation of the occupation numbers for a finite N , as shown in figure 1, and in turn yields the oscillatory behavior of the inverse participation ratio. Note that the period of oscillation is p , as one may expect.
- [18] R. S. Berry in *Structure and Dynamics of Atoms and Molecules: Conceptual Trends*, edited by J. L. Calais and E. S. Kryachko (Kluwer, Dordrecht, 1995) p. 155, and references therein.
- [19] The difference between CPD and the usual boson density can be interpreted as follows: the CPD describes bosons in their *intrinsic (body-fixed)* frame of reference, while the density distribution describes bosons in the laboratory frame of reference where the rotational and center-of-mass displacements are superimposed upon the intrinsic probability density. Note that the CPD can be rewritten as: $\mathcal{P}(\mathbf{r} | \mathbf{v}_0) = \rho(\mathbf{r})g^{(2)}(\mathbf{r}, \mathbf{v}_0)$, where $\rho(\mathbf{r})$ and $g^{(2)}(\mathbf{r}, \mathbf{v}_0)$ are the density distribution and normalized second-order correlation function, respectively. For a single condensate, $g^{(2)}(\mathbf{r}, \mathbf{v}_0) = 1$, and the CPD is simply reduced to the boson density, i.e., $\mathcal{P}(\mathbf{r} | \mathbf{v}_0) = \rho(\mathbf{r})$.
- [20] A. J. Leggett and F. Sols, *Found. Phys.* **21**, 353 (1991).

Figures Captions

Fig. 1. (color) Values of n_j (in blue) and Δn_j (in red) of five lowest SP oscillator states as a function of L for a system of $N = 40$ bosons with $j_{\max} = 6$. The cases of $j = 5$ and 6 are not shown due to their low occupancy.

Fig. 2. $1/I_C$ versus L/N for a system of $N = 30, 40, 50$, and 60 bosons. Small oscillations at $L/N > 1.6$ are caused by the finite N effect. The overall profile of $1/I_C$ is nearly independent of N , thus $1/I_C$ can be used to qualitatively measure the number of macroscopic occupation numbers in the limit of $N \rightarrow \infty$.

Fig. 3. (color) V_{int} in units of gN^2 as a function of L/N for a system of $N = 20$ (in green), 40 (in blue), and 60 (in red) bosons. For comparison, the V_{int} predicted by MFT is also depicted by the dark solid line. The inset shows the energy difference $\Delta V_{int} = V_{int}^{mf} - V_{int}$ in units of gN . Note that all the ΔV_{int} with different N are approximately located on a universal curve.

Fig. 4. Selected CPDs for a system of $N = 40$ bosons. (a), (b), (c) and (d) correspond to the emergence of vortex-like patterns with p -fold symmetry ($p = 1, 2, 3, 4$). In each panel, L increases in steps of one unit, and the starting value of L in (a), (b), (c), and (d) is 33, 62, 79, and 108, respectively. The values of x and y in each subplot range from -3.0 to $+3.0$. The given point \mathbf{v}_0 is $(0, 1.0)$. For large N , the CPD is nearly independent of \mathbf{v}_0 .




Fig. 1 (color)
 Xiaji Liu *et al.*
 Fragmented condensate ...




Fig. 2
 Xiaji Liu *et al.*
 Fragmented condensate ...




Fig. 3 (color)
Xiaji Liu *et al.*
Fragmented condensate ...